



Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	July 9, 2021 August 4, 2021

ADUHELM™ (aducanumab-avwa)

LENGTH OF AUTHORIZATION: 6 months

REVIEW CRITERIA:

- Patient must be ≥ 18 years of age; **AND**
- Patient has mild cognitive impairment (MCI) due to Alzheimer's disease or mild Alzheimer's dementia (there is insufficient evidence in moderate or severe AD) as evidenced by all the following:
 - Clinical Dementia Rating (CDR)-Global Score of 0.5; **AND**
 - Objective evidence of cognitive impairment at screening; **AND**
 - Mini-Mental Status Exam (MMSE) score between 24 and 30 (inclusive); **AND**
 - Positron Emission Tomography (PET) scan is positive for amyloid beta plaque; **AND**
- Other conditions mimicking, but of non-Alzheimer's dementia etiology, have been ruled out (e.g., vascular dementia, dementia with Lewy bodies [DLB], frontotemporal dementia [FTD], normal pressure hydrocephalus); **AND**
- Drug must be prescribed by, or in consultation with, a specialist in neurology or gerontology; **AND**
- Patient has received a baseline brain magnetic resonance imaging (MRI) prior to initiating treatment (within 1 year prior); **AND**
- Patient does not have any of the following within 1 year of treatment initiation: pre-treatment localized superficial siderosis, ≥ 10 brain microhemorrhages, or brain hemorrhage > 1 cm; **AND**
- Prescriber has assessed and documented baseline disease severity utilizing an objective measure/tool (e.g., MMSE, Alzheimer's Disease Assessment Scale-Cognitive Subscale [ADAS-Cog-13], Alzheimer's Disease Cooperative Study-Activities of Daily Living Inventory-Mild Cognitive Impairment version [ADCS-ADL-MCI], Clinical Dementia Rating-Sum of Boxes [CDR-SB]).

CONTINUATION OF THERAPY:

- Patient continues to meet the above criteria; **AND**
- Patient has not had unacceptable toxicity from the drug (e.g., amyloid related imaging abnormalities [ARIA]-edema [ARIA-E], severe hypersensitivity reactions); **AND**
- Patient has responded to therapy compared to pre-treatment baseline as evidenced by improvement, stability, or slowing in cognitive and/or functional impairment in ≥ 1 of the following (not all-inclusive) objective measures assessed and documented at baseline: ADAS-Cog 13, ADCS-ADL-MCI, MMSE, or CDR-SB; **AND**
- Patient has not progressed to moderate or severe AD; **AND**
- Patient must continue maintenance therapy at the recommended dosage per product labeling; **AND**



Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	July 9, 2021 August 4, 2021

- Patient has received MRI monitoring throughout therapy including an MRI prior to the 7th and 12th doses for monitoring of ARIA-hemosiderin (ARIA-H) microhemorrhages; **AND**
- Patient has < 10 new incident microhemorrhages **and** ≤ 2 focal areas of superficial siderosis (radiographic mild to moderate ARIA-H) observed; **OR**
- Patient has ≥ 10 new incident microhemorrhages or > 2 focal areas of superficial siderosis (radiographic severe ARIA-H[†]) are observed and patient meets the following criteria:
 - Treatment is continued with caution only after a clinical evaluation; **AND**
 - Subsequent follow-up MRI demonstrates radiographic stabilization (e.g., no increase in size or number of ARIA-H[†]).

ARIA Classification and Radiographic Severity [†]			
Severity	ARIA-E (based on FLAIR hyperintensity)	ARIA-H microhemorrhage (quantity of new incident microhemorrhages)	ARIA-H superficial siderosis (quantity of superficial siderosis focal areas)
Mild	confined to sulcus and or cortex/subcortical white matter in 1 location < 5 cm	≤ 4	1
Moderate	5 to 10 cm, or > 1 site of involvement, each measuring < 10 cm	5 to 9	2
Severe	> 10 cm, often with significant subcortical white matter and/or sulcal involvement; ≥ 1 separate sites of involvement may be noted	≥ 10	>2

DOSING AND ADMINISTRATION:

- Refer to product labeling at <https://www.biogen.com/us/aduhelm-pi.pdf>
- Available as 170 mg/1.7 mL and 300 mg/3 mL solution in a single-dose vial for intravenous infusion