



Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	March 9, 2021 October 4, 2021, March 31, 2022

## **AMONDYS 45 (casimersen)**

**LENGTH OF AUTHORIZATION:** SIX MONTHS

### **REVIEW CRITERIA:**

- Patient must have the diagnosis of Duchenne muscular dystrophy (DMD).
- Submission of medical records (e.g., chart notes, laboratory values) as genetic test is required to confirm that a patient's mutation of the DMD gene is amenable to exon 45 skipping.
- Medication is prescribed by or in consultation with a neurologist or a physician who specializes in treatment of DMD (i.e., pediatric neurologist, cardiologist or pulmonary specialist).
- Submission of cystatin C, urine dipstick, and urine protein-to-creatinine ratio prior to starting therapy.
- Patient has been on stable dose of oral corticosteroids for at least 24 weeks prior to starting therapy **unless contraindicated or intolerant**.
- Patient is not concurrently treated with other DMD antisense oligonucleotides (e.g., Golodirsen, Viltolarsen, or Eteplirsen).
- If the patient is ambulatory, functional level determination of baseline assessment of ambulatory function (six-minute walk test (6MWT), time to run/walk 10-meter test (TTRW), time to climb 4-stair test (TTCLIMB), time to stand (TTSTAND) or North Star Ambulatory Assessment (NSAA)) is required.
- If not ambulatory, patient must have a Brooke Upper Extremity Function Scale of five or less documented OR a Forced Vital Capacity of 30% or more.

### **CONTINUATION OF THERAPY:**

- Patient met initial review criteria.
- Documentation of improvement, maintenance, or slowing of disease progression:
  - For ambulatory patients – submission of 6MWT, TTRW, TTCLIMB, TTSTAND or NSAA.
  - For non-ambulatory patients – submission of Brooke Upper Extremity Function Scale (five or less) documented OR a Forced Vital Capacity document (30% or more).
- Submission of cystatin C, urine dipstick, and urine protein-to-creatinine ratio.
- Patient is not concurrently treated with other DMD antisense oligonucleotides (e.g., Golodirsen, Viltolarsen, or Eteplirsen).

### **DOSING AND ADMINISTRATION:**

- Refer to product labeling at <https://www.accessdata.fda.gov/scripts/cder/daf/>
- Available as 100 mg/2 mL single-dose vial