



Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	January 7, 2022 August 5, 2022, October 14, 2022

Thrombopoiesis Stimulating Agents

Non-Preferred Agents: Doptelet[®] (avatrombopag), Mulpleta[®] (lusutrombopag), Nplate[®] (romiplostim), Promacta[®] (eltrombopag), Tavalisse[™] (fostamatinib)

LENGTH OF AUTHORIZATION: Up to 6 months for *ITP, Severe Aplastic Anemia and HS-ARS*
Up to 4 months for *Chronic Hepatitis C-associated Thrombocytopenia*
Up to 1 week for *Thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure*

INITIAL REVIEW CRITERIA:

- Diagnosis confirmed by supporting documentation.

Chronic immune (idiopathic) thrombocytopenia (ITP) - Avatrombopag (Doptelet[®]), Eltrombopag (Promacta[®]) or Romiplostim (Nplate[®])

- Documentation of platelet count less than $50 \times 10^9/L$ ($50,000/mm^3$) and/or signs and symptoms of a low platelet count (bruising, petechiae, bleeding from nostrils, gums, etc.).
- Patient has history of failure, intolerance or contraindication to **ONE** of the following:
 - Glucocorticoids
 - Intravenous immune globulin (IVIG)
 - Rituximab
 - History of splenectomy

Severe aplastic anemia - Eltrombopag (Promacta[®])

- Patient is ≥ 2 years of age.
- Lab documentation of the following:
 - Platelet count of less than $30 \times 10^9/L$ ($30,000/mm^3$) **or** patient is platelet transfusion dependent.
 - Hemoglobin 8.4 g/dL or lower **or** patient is dependent on transfusions of red blood cells (RBCs)
 - Absolute neutrophil count (ANC) approximating $0.5 \times 10^9/L$.
- Promacta[®] must be used in combination with standard immunosuppressive therapy. History of failure, intolerance or contraindication must be documented.

Chronic Hepatitis C-associated Thrombocytopenia - Eltrombopag (Promacta[®])

- Documentation the hepatitis C treatment regimen is used in conjunction with interferon-based therapy.
- Documentation of platelet count less than $50 \times 10^9/L$ ($50,000/mm^3$).

Thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure - Avatrombopag (Doptelet[®]) or Lusutrombopag (Mulpleta[®])

- Patient is ≥ 18 years of age.
- Documentation of platelet count less than $50 \times 10^9/L$ ($50,000/mm^3$)



Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	January 7, 2022 August 5, 2022, October 14, 2022

Hematopoietic Syndrome of Acute Radiation Syndrome [HS-ARS] - Romiplostim (Nplate®)

- Patient receiving myelosuppressive doses of radiation

CONTINUATION OF THERAPY REVIEW CRITERIA:

Chronic immune (idiopathic) thrombocytopenia (ITP) - Avatrombopag (Doptelet®), Eltrombopag (Promacta®) or Romiplostim (Nplate®)

- Documentation of platelet count greater than $50 \times 10^9/L$ ($50,000/mm^3$)

Severe aplastic anemia - Eltrombopag (Promacta®)

- Platelet count increases to $20 \times 10^9/L$ above baseline, **or** stable platelet counts with transfusion independence for a minimum of 8 weeks.
- Hemoglobin increase by greater than 1.5 g/dL or a reduction in greater than or equal to 4 units of RBC transfusions for 8 consecutive weeks.
- ANC increase of 100% or an ANC increase greater than $0.5 \times 10^9 /L$.

***If the patient has not met at least one of the above criteria after 16 weeks of treatment, continuation of therapy should NOT be approved.**

DOSING AND ADMINISTRATION:

- Refer to product labeling at <https://www.accessdata.fda.gov/scripts/cder/daf/>